arXiv:1202.4130v1 [hep-lat] 19 Feb 2012

Gauge-Invariant Formalism with Dirac-mode Expansion for Confinement and Chiral Symmetry Breaking

Shinya Gongyo,[*] Takumi Iritani,[†] and Hideo Suganuma[‡]
Department of Physics, Graduate School of Science, Kyoto University,
Kitashirakawa-oiwake, Sakyo, Kyoto 606-8502, Japan

(Dated: April 5, 2012)

We develop a manifestly gauge-covariant expansion of the QCD operator such as the Wilson loop, using the eigen-mode of the QCD Dirac operator $\not{D} = \gamma^\mu D^\mu$. With this method, we perform a direct analysis of the correlation between confinement and chiral symmetry breaking in lattice QCD Monte Carlo calculation on 6^4 at β=5.6. As a remarkable fact, the confinement force is almost unchanged even after removing the low-lying Dirac modes, which are responsible to chiral symmetry breaking. This indicates that one-to-one correspondence does not hold for between confinement and chiral symmetry breaking in QCD. In this analysis, we carefully amputate only the "essence of chiral symmetry breaking" by cutting off the low-lying Dirac modes, and can artificially realize the "confined but chiral restored situation" in QCD.

PACS numbers: 12.38.Aw, 12.38.Gc, 14.70.Dj

I. INTRODUCTION

Nowadays, quantum chromodynamics (QCD) has been established as the fundamental gauge theory of the strong interaction. However, nonperturbative properties of low-energy QCD such as color confinement and chiral symmetry breaking [1] are not yet well understood, which gives one of the most difficult problems in theoretical physics. The nonperturbative QCD has been studied in lattice QCD [2–6] and various analytical frameworks [7–12].

In particular, it is rather interesting and important to examine the correlation between confinement and chiral symmetry breaking [12–18], since the direct relation is not yet shown between them in QCD. The strong correlation between them has been suggested by the almost simultaneous phase transitions of deconfinement and chiral restoration in lattice QCD both at finite temperature [5, 19] and in a small-volume box [5].

The close relation between confinement and chiral symmetry breaking has been also suggested in terms of the monopole degrees of freedom [12–14]. Here, the monopole topologically appears in QCD by taking the maximally Abelian (MA) gauge [20–24]. For example, by removing the monopoles in the MA gauge, confinement and chiral symmetry breaking are simultaneously lost in lattice QCD [13, 14]. (The instantons also disappear without monopoles [22].) This indicates an important role of the monopole to both confinement and chiral symmetry breaking, and these two nonperturbative QCD phenomena seem to be related via the monopole.

However, as a possibility, removing the monopoles may be "too fatal" for most nonperturbative properties. If this is the case, nonperturbative QCD phenomena are simultaneously lost by their cut.

In fact, *if only the relevant ingredient of chiral symmetry breaking is carefully removed, how will be confinement?*

In this paper, to get the answer, we perform a direct investigation between color confinement and chiral symmetry breaking in lattice QCD, using the Dirac-mode expansion in a gauge-invariant manner [25].

II. GAUGE-INVARIANT FORMALISM WITH DIRAC-MODE EXPANSION

We newly develop a manifestly gauge-covariant expansion of the QCD operator such as the Wilson loop, using the eigen-mode of the QCD Dirac operator $\not{D} = \gamma^\mu D^\mu$, and investigate the relation between confinement and chiral symmetry breaking.

A. Gauge Covariant Expansion in QCD instead of Fourier Expansion

In the previous studies [26, 27], we investigated the relevant gluon-momentum region for confinement in lattice QCD, and found that the string tension σ, i.e., the confining force, is almost unchanged even after removing the high-momentum gluon component above 1.5GeV in the Landau gauge. In fact, the confinement property originates from the low-momentum gluon component below 1.5GeV, which is the upper limit to contribute to σ.

The previous study on the relevant gluonic modes was based on the Fourier expansion, i.e., the eigen-mode expansion of the momentum operator p^μ. Because of the commutable nature of $[p^\mu, p^\nu] = 0$, all the momentum p^μ can be simultaneously diagonalized. which is one of the strong merits of the Fourier expansion. Also it keeps Lorentz covariance.

---

[*] Electronic address: gongyo@ruby.scphys.kyoto-u.ac.jp
[†] Electronic address: iritani@ruby.scphys.kyoto-u.ac.jp
[‡] Electronic address: suganuma@ruby.scphys.kyoto-u.ac.jp

However, the Fourier expansion does *not* keep gauge invariance in gauge theories. Therefore, for the use of the Fourier expansion in QCD, one has to select a suitable gauge such as the Landau gauge [26–28], where the gauge-field fluctuation is strongly suppressed in Euclidean QCD.

As a next challenge, we consider a gauge-invariant method, using a gauge-covariant expansion in QCD instead of the Fourier expansion. In fact, we consider a generalization of the Fourier expansion or an alternative expansion with keeping the gauge symmetry.

A straight generalization is to use the covariant derivative operator D^μ instead of the derivative operator ∂^μ. However, due to the non-commutable nature of $[D^\mu, D^\nu] \neq 0$, one cannot diagonalize all the covariant derivative D^μ $(\mu = 1, 2, 3, 4)$ simultaneously, but only one of them can be diagonalized. For example, the eigenmode expansion of D_4 keeps gauge covariance and is rather interesting, but this type of the expansion inevitably breaks the Lorentz covariance. Then, we consider the eigen-mode expansion of the Dirac operator $\not{D} = \gamma^\mu D^\mu$ or $D^2 = D^\mu D^\mu$, since such an expansion keeps both gauge symmetry and Lorentz covariance.

In particular, the Dirac-mode expansion is rather interesting because the Dirac operator $\not{D}$ directly connects with chiral symmetry breaking via the Banks-Casher relation [8] and its zero modes are related to the topological charge via the Atiyah-Singer index theorem [29]. Here, we mainly consider the manifestly gauge-invariant new method using the Dirac-mode expansion. Thus, the Dirac-mode expansion has some important merits.

- The Dirac-mode expansion method manifestly keeps both gauge and Lorentz invariance.
- Each QCD phenomenon can be directly investigated in terms of chiral symmetry breaking.

B. Eigen-mode of Dirac Operator in Lattice QCD

Now, we consider the Dirac operator and its eigenmodes in lattice QCD formalism with spacing a in the Euclidean metric. On the lattice, each site is labeled by $x = (x_1, x_2, x_3, x_4)$ with x_μ being an integer. In lattice QCD, the gauge field is described by the link-variable $U_\mu(x) = e^{iagA_\mu(x)} \in \mathrm{SU}(N_c)$, where g is the QCD gauge coupling and $A_\mu(x) \in \mathrm{su}(N_c)$ corresponds to the gluon field.

In lattice QCD, the Dirac operator $\not{D} = \gamma_\mu D_\mu$ is expressed with $U_\mu(x)$ as

$$\not{D}_{x,y} \equiv \frac{1}{2a}\sum_{\mu=1}^{4} \gamma_\mu \left[U_\mu(x)\delta_{x+\hat{\mu},y} - U_{-\mu}(x)\delta_{x-\hat{\mu},y}\right], \quad (1)$$

where the convenient notation $U_{-\mu}(x) \equiv U^\dagger_\mu(x-\hat{\mu})$ is used. Here, $\hat{\mu}$ denotes the unit vector on the lattice in μ-direction [5].

In this paper, we adopt the hermite definition of the γ-matrix, $\gamma^\dagger_\mu = \gamma_\mu$. Then, $\not{D}$ is anti-hermite and satisfies

$$\not{D}^\dagger_{y,x} = -\not{D}_{x,y}. \quad (2)$$

The normalized eigen-state $|n\rangle$ of the Dirac operator $\not{D}$ is introduced as

$$\not{D}|n\rangle = i\lambda_n|n\rangle \quad (3)$$

with $\lambda_n \in \mathbf{R}$. Because of $\{\gamma_5, \not{D}\} = 0$, the state $\gamma_5|n\rangle$ is also an eigen-state of $\not{D}$ with the eigenvalue $-i\lambda_n$. The Dirac eigenfunction

$$\psi_n(x) \equiv \langle x|n\rangle \quad (4)$$

obeys $\not{D}\psi_n(x) = i\lambda_n\psi_n(x)$, and its explicit form of the eigenvalue equation in lattice QCD is

$$\begin{aligned} &\tfrac{1}{2a}\sum_{\mu=1}^{4}\gamma_\mu[U_\mu(x)\psi_n(x+\hat{\mu}) - U_{-\mu}(x)\psi_n(x-\hat{\mu})] \\ &= i\lambda_n\psi_n(x). \end{aligned} \quad (5)$$

The Dirac eigenfunction $\psi_n(x)$ can be numerically obtained in lattice QCD, besides a phase factor.

According to $U_\mu(x) \to V(x)U_\mu(x)V^\dagger(x+\hat{\mu})$, the gauge transformation of $\psi_n(x)$ is found to be

$$\psi_n(x) \to V(x)\psi_n(x), \quad (6)$$

which is the same as that of the quark field. To be strict, for the Dirac eigenfunction, there appears an irrelevant n-dependent global phase factor $e^{i\varphi_n[V]}$, according to the arbitrariness of the definition of $\psi_n(x)$.

It is notable that the quark condensate $\langle\bar{q}q\rangle$, the order parameter of chiral symmetry breaking, is given by the zero-eigenvalue density $\rho(0)$ of the Dirac operator, via the Banks-Casher relation [8],

$$\langle\bar{q}q\rangle = -\lim_{m\to 0}\lim_{V\to\infty}\pi\rho(0). \quad (7)$$

Here, the spectral density of the Dirac operator is defined by

$$\rho(\lambda) \equiv \frac{1}{V}\sum_n\langle\delta(\lambda-\lambda_n)\rangle, \quad (8)$$

with the four-dimensional volume V. Also, the zero-mode number asymmetry of the Dirac operator $\not{D}$ is equal to the topological charge (the instanton number) $Q \equiv \frac{g^2}{16\pi^2}\int d^4x \ \mathrm{Tr}\ (G_{\mu\nu}\tilde{G}_{\mu\nu})$, which is known as the Atiyah-Singer index theorem, Index($\not{D}$)=Q [29].

C. Kogut-Susskind Technique to reduce Dirac Eigenvalue Calculation

In calculating the eigenvalue of the Dirac operator $\not{D}$, we use the Kogut-Susskind (KS) technique [3, 5], which is often used to remove the redundant doublers of lattice fermions. Here, the use of the KS technique is just the practical reason to reduce the calculation of the Dirac eigenvalues. In fact, the result of the Dirac-mode projection, which will be shown in Sec.IV, is unchanged, when the Dirac operator is directly diagonalized.

In the KS technique, one introduces

$$T(x) \equiv \gamma_1^{x_1}\gamma_2^{x_2}\gamma_3^{x_3}\gamma_4^{x_4}, \qquad (9)$$

which satisfies $T(x-\hat{\mu}) = T(x+\hat{\mu})$, due to $\gamma_\mu^2 = 1$ for each μ. As a key relation, "diagonalization" of all the gamma matrices γ_μ can be done as

$$T(x)^\dagger \gamma_\mu T(x+\hat{\mu}) = \eta_\mu(x) 1 \qquad (10)$$

with the staggered phase $\eta_\mu(x)$ defined by

$$\eta_1(x) \equiv 1, \quad \eta_\mu(x) = (-1)^{x_1+\cdots+x_{\mu-1}} \ (\mu \geq 2). \qquad (11)$$

By introducing $\chi_n(x) \equiv T^\dagger(x)\psi_n(x)$, the eigenvalue equation is reduced into

$$\begin{aligned} \tfrac{1}{2a} \sum_{\mu=1}^{4} &\eta_\mu(x)[U_\mu(x)\chi_n(x+\hat{\mu}) - U_{-\mu}(x)\chi_n(x-\hat{\mu})] \\ = &\ i\lambda_n \chi_n(x), \qquad (12) \end{aligned}$$

which does not include spinor indices. In the KS method, the Dirac operator $\gamma_\mu D_\mu$ is replaced by the KS Dirac operator $\eta_\mu D_\mu$ with dropping the spinor degrees of freedom, which reduces the lattice-fermion species from 16 to 4 [5].

To summarize the KS method, there is no spin indices, and the explicit form of the KS Dirac operator $\eta_\mu D_\mu$ is

$$(\eta_\mu D_\mu)_{x,y} = \frac{1}{2a}\sum_{\mu=1}^{4} \eta_\mu(x)[U_\mu(x)\delta_{x+\hat{\mu},y} - U_{-\mu}(x)\delta_{x-\hat{\mu},y}]. \qquad (13)$$

The spinless eigenfunction $\chi_n(x)$ satisfies Eq.(12), i.e.,

$$\eta_\mu D_\mu \chi_n(x) \equiv \sum_y \sum_{\mu=1}^{4} (\eta_\mu D_\mu)_{x,y}\chi_n(y) = i\lambda_n \chi_n(x). \qquad (14)$$

In the KS method, the chiral partner $\gamma_5\psi_n(x)$ reduces into $(-1)^{x_1+x_2+x_3+x_4}\chi_n(x)$, which is an eigenfunction of $\eta_\mu D_\mu$ with the eigenvalue $-i\lambda_n$.

Using the KS technique [3, 5], the Dirac-mode number $L^4 \times N_c \times 4$ is reduced to be $L^4 \times N_c$ on the L^4 lattice. The actual number of the independent Dirac eigenvalue λ_n is about $L^4 \times N_c/2$, due to the chiral property of the Dirac operator, i.e., pairwise appearance of $\pm\lambda_n$, apart from chiral zero modes.

III. OPERATOR FORMALISM IN LATTICE QCD

To keep the gauge symmetry, careful treatments are necessary, since naive approximations may break the gauge symmetry. Here, we take the "operator formalism" [25], as explained below.

We define the link-variable operator $\hat{U}_{\pm\mu}$ by the matrix element of

$$\langle x|\hat{U}_{\pm\mu}|y\rangle = U_{\pm\mu}(x)\delta_{x\pm\hat{\mu},y}. \qquad (15)$$

Note that $\hat{U}_\mu$ and $\hat{U}_{-\mu}$ are hermite conjugate as the operator in the Hilbert space in the sense that

$$\begin{aligned} \langle y|\hat{U}_\mu^\dagger|x\rangle &= U_\mu^\dagger(y)\delta_{y+\hat{\mu},x} = U_\mu^\dagger(x-\hat{\mu})\delta_{x-\hat{\mu},y} \\ &= U_{-\mu}(x)\delta_{x-\hat{\mu},y} = \langle x|\hat{U}_{-\mu}|y\rangle. \qquad (16) \end{aligned}$$

In the operator formalism, Eq.(5) for the Dirac eigenstate is simply expressed as

$$\frac{1}{2a}\sum_{\mu=1}^{4} \gamma_\mu(\hat{U}_\mu - \hat{U}_{-\mu})|n\rangle = i\lambda_n|n\rangle. \qquad (17)$$

In the KS method, where the spinor degrees of freedom is dropped off, one identifies $\chi_n(x) = \langle x|n\rangle$, and then Eq.(12) for the KS Dirac eigen-state is expressed as

$$\frac{1}{2a}\sum_{\mu=1}^{4} \hat{\eta}_\mu(\hat{U}_\mu - \hat{U}_{-\mu})|n\rangle = i\lambda_n|n\rangle, \qquad (18)$$

where $\hat{\eta}_\mu$ is defined by the matrix element of $\langle x|\hat{\eta}_\mu|y\rangle = \eta_\mu(x)\delta_{x,y}$. Owing to $\eta_\mu(x \pm \hat{\mu}) = \eta_\mu(x)$, one finds $\hat{\eta}_\mu\hat{U}_{\pm\mu} = \hat{U}_{\pm\mu}\hat{\eta}_\mu$, so that there is no ordering uncertainty in the KS Dirac operator in Eq.(18).

In the following, we mainly use the ordinary Dirac operator $\gamma_\mu D_\mu$ and the spinor eigenfunction $\psi_n(x) = \langle x|n\rangle$. When the KS method is applied, one only has to use the identification of $\chi_n(x) = \langle x|n\rangle$ in the following arguments. The final results are the same between both calculations based on $\gamma_\mu D_\mu$ and $\eta_\mu D_\mu$.

The Wilson-loop operator $\hat{W}$ is defined as the product of $\hat{U}_\mu$ along a rectangular loop,

$$\hat{W} \equiv \prod_{k=1}^{N} \hat{U}_{\mu_k} = \hat{U}_{\mu_1}\hat{U}_{\mu_2}\cdots\hat{U}_{\mu_N}. \qquad (19)$$

For arbitrary loops, one finds $\sum_{k=1}^{N}\hat{\mu}_k = 0$. We note that the functional trace of the Wilson-loop operator $\hat{W}$ is proportional to the ordinary vacuum expectation value $\langle W\rangle$ of the Wilson loop:

$$
\begin{aligned}
\mathrm{Tr}\ \hat{W} &= \mathrm{tr}\sum_x \langle x|\hat{W}|x\rangle = \mathrm{tr}\sum_x \langle x|\hat{U}_{\mu_1}\hat{U}_{\mu_2}\cdots\hat{U}_{\mu_N}|x\rangle \\
&= \mathrm{tr}\sum_{x_1,x_2,\cdots,x_N} \langle x_1|\hat{U}_{\mu_1}|x_2\rangle\langle x_2|\hat{U}_{\mu_2}|x_3\rangle\langle x_3|\hat{U}_{\mu_3}|x_4\rangle\cdots\langle x_N|\hat{U}_{\mu_N}|x_1\rangle \\
&= \mathrm{tr}\sum_x \langle x|\hat{U}_{\mu_1}|x+\hat{\mu}_1\rangle\langle x+\hat{\mu}_1|\hat{U}_{\mu_2}|x+\sum_{k=1}^{2}\hat{\mu}_k\rangle\cdots\langle x+\sum_{k=1}^{N}|\hat{U}_{\mu_N}|x\rangle \\
&= \sum_x \mathrm{tr}\{U_{\mu_1}(x)U_{\mu_2}(x+\hat{\mu}_1)U_{\mu_3}(x+\sum_{k=1}^{2}\hat{\mu}_k)\cdots U_{\mu_N}(x+\sum_{k=1}^{N}\hat{\mu}_k)\} \\
&= \langle W\rangle\cdot \mathrm{Tr}\ 1.
\end{aligned}
\tag{20}
$$

Here, "Tr" denotes the functional trace, and "tr" the trace over SU(3) color index.

The Dirac-mode matrix element of the link-variable operator $\hat{U}_\mu$ can be expressed with $\psi_n(x)$:

$$
\begin{aligned}
\langle m|\hat{U}|n\rangle &= \sum_x \langle m|x\rangle\langle x|\hat{U}_\mu|x+\hat{\mu}\rangle\langle x+\hat{\mu}|n\rangle \\
&= \sum_x \psi_m^\dagger(x)U_\mu(x)\psi_n(x+\hat{\mu}).
\end{aligned}
\tag{21}
$$

Although the total number of the matrix element is very huge, the matrix element is calculable and gauge invariant, apart from an irrelevant phase factor. Using the gauge transformation (6), we find the gauge transformation of the matrix element as

$$
\begin{aligned}
\langle m|\hat{U}_\mu|n\rangle &= \sum_x \psi_m^\dagger(x)U_\mu(x)\psi_n(x+\hat{\mu}) \\
&\to \sum_x \psi_m^\dagger(x)V^\dagger(x)\cdot V(x)U_\mu(x)V^\dagger(x+\hat{\mu})\cdot V(x+\hat{\mu})\psi_n(x+\hat{\mu}) \\
&= \sum_x \psi_m^\dagger(x)U_\mu(x)\psi_n(x+\hat{\mu}) = \langle m|\hat{U}_\mu|n\rangle.
\end{aligned}
\tag{22}
$$

To be strict, there appears an n-dependent global phase factor, corresponding to the arbitrariness of the phase in the basis $|n\rangle$. However, this phase factor cancels as $e^{-i\varphi_n}e^{i\varphi_n}=1$ between $|n\rangle$ and $\langle n|$, and does not appear for QCD physical quantities including the Wilson loop.

In the practical lattice-QCD calculation, we adopt the KS technique to reduce the computational complexity, as mentioned in Sec II-C. In the KS method, instead of $\psi_n(x)$, we use the spinless eigenfunction $\chi_n(x)$ of the KS Dirac operator $\eta_\mu D_\mu$, with the identification of $\chi_n(x)=\langle x|n\rangle$, and the KS-reduced matrix element of $\hat{U}_\mu$ is expressed as

$$
\begin{aligned}
\langle m|\hat{U}|n\rangle &= \sum_x \langle m|x\rangle\langle x|\hat{U}_\mu|x+\hat{\mu}\rangle\langle x+\hat{\mu}|n\rangle \\
&= \sum_x \chi_m^\dagger(x)U_\mu(x)\chi_n(x+\hat{\mu}).
\end{aligned}
\tag{23}
$$

In the arguments in the next section, the same results are obtained between the calculations based on the original Dirac operator $\gamma_\mu D_\mu$ and the KS Dirac operator $\eta_\mu D_\mu$.

IV. DIRAC-MODE EXPANSION AND PROJECTION

From the completeness of the Dirac-mode basis, $\sum_n |n\rangle\langle n|=1$, we get

$$\hat{O}=\sum_m\sum_n |m\rangle\langle m|\hat{O}|n\rangle\langle n| \tag{24}$$

for arbitrary operators. Based on this relation, the Dirac-mode expansion and projection can be defined [25]. We define the projection operator $\hat{P}$ which restricts the Dirac-mode space,

$$\hat{P}\equiv\sum_{n\in A}|n\rangle\langle n|, \tag{25}$$

where A denotes arbitrary set of Dirac modes. In $\hat{P}$, the arbitrary phase cancels between $|n\rangle$ and $\langle n|$. One finds

$\hat{P}^2=\hat{P}$ and $\hat{P}^\dagger=\hat{P}$. The typical projections are IR-cut and UV-cut of the Dirac modes:

$$\hat{P}_{\rm IR}\equiv\sum_{|\lambda_n|\ge\Lambda_{\rm IR}}|n\rangle\langle n|,\quad \hat{P}_{\rm UV}\equiv\sum_{|\lambda_n|\le\Lambda_{\rm UV}}|n\rangle\langle n|. \qquad (26)$$

Using the projection operator $\hat{P}$, we define the Dirac-mode projected link-variable operator,

$$\hat{U}_\mu^P\equiv\hat{P}\hat{U}_\mu\hat{P}=\sum_{m\in A}\sum_{n\in A}|m\rangle\langle m|\hat{U}_\mu|n\rangle\langle n|. \qquad (27)$$

During this projection, there appears some nonlocality in general, but it would not be important for the argument of large-distance properties such as confinement. From the Wilson-loop operator

$$\hat{W}\equiv\prod_{k=1}^{N}\hat{U}_{\mu_k}, \qquad (28)$$

we define the Dirac-mode projected Wilson-loop operator,

$$\begin{aligned}\hat{W}^P &\equiv \prod_{k=1}^{N}\hat{U}_{\mu_k}^P=\hat{U}_{\mu_1}^P\hat{U}_{\mu_2}^P\cdots\hat{U}_{\mu_N}^P=\hat{P}\hat{U}_{\mu_1}\hat{P}\hat{U}_{\mu_2}\hat{P}\cdots\hat{P}\hat{U}_{\mu_N}\hat{P}\\ &= \sum_{n_1,n_2,\cdots,n_{N+1}\in A}|n_1\rangle\langle n_1|\hat{U}_{\mu_1}|n_2\rangle\langle n_2|\hat{U}_{\mu_2}|n_3\rangle\cdots\langle n_N|\hat{U}_{\mu_N}|n_{N+1}\rangle\langle n_{N+1}|. \end{aligned}\qquad (29)$$

Then, we obtain the functional trace of the Dirac-mode projected Wilson-loop operator,

$$\begin{aligned}{\rm Tr}\ \hat{W}^P &= {\rm Tr}\ \prod_{k=1}^{N}\hat{U}_{\mu_k}^P={\rm Tr}\ \hat{U}_{\mu_1}^P\hat{U}_{\mu_2}^P\cdots\hat{U}_{\mu_N}^P={\rm Tr}\ \hat{P}\hat{U}_{\mu_1}\hat{P}\hat{U}_{\mu_2}\hat{P}\cdots\hat{P}\hat{U}_{\mu_N}\hat{P}\\ &= {\rm tr}\sum_{n_1,n_2,\cdots,n_N\in A}\langle n_1|\hat{U}_{\mu_1}|n_2\rangle\langle n_2|\hat{U}_{\mu_2}|n_3\rangle\cdots\langle n_N|\hat{U}_{\mu_N}|n_1\rangle, \end{aligned}\qquad (30)$$

which is manifestly gauge invariant. Here, the arbitrary phase factor cancels between $|n_k\rangle$ and $\langle n_k|$. Its gauge invariance is also numerically checked in the lattice QCD Monte Carlo calculation.

From Tr $\hat{W}^P(R,T)$ corresponding to the $R\times T$ rectangular loop, we define the Dirac-mode projected inter-quark potential,

$$V^P(R)\equiv-\lim_{T\to\infty}\frac{1}{T}\ln\{{\rm Tr}\ \hat{W}^P(R,T)\}, \qquad (31)$$

which is also manifestly gauge-invariant. In the unprojected case of $\hat{P}=1$, the ordinary inter-quark potential is obtained apart from an irrelevant constant,

$$\begin{aligned}V(R) &= -\lim_{T\to\infty}\frac{1}{T}\ln\{{\rm Tr}\ \hat{W}(R,T)\}\\ &= -\lim_{T\to\infty}\frac{1}{T}\ln\langle W(R,T)\rangle+\text{irrelevant const.}, \end{aligned}\qquad (32)$$

because of Tr $\hat{W}=\langle W\rangle\cdot{\rm Tr}\ 1$, as was derived in Eq.(20).

V. ANALYSIS OF CONFINEMENT IN TERMS OF DIRAC MODES IN QCD

We consider various projection space A in the Dirac-mode space, e.g., IR-cut or UV-cut of Dirac modes. With this Dirac-mode expansion and projection formalism, we calculate the Dirac-mode projected inter-quark potential $V^P(R)$ in a gauge-invariant manner. In particular, using IR-cut of the Dirac modes, we directly investigate the role of low-lying Dirac modes to confinement, since the low-lying modes are responsible to chiral symmetry breaking.

As a technical difficulty of this formalism, we have to deal with huge dimensional matrices and their products. Actually, the total matrix dimension of $\langle m|\hat{U}_\mu|n\rangle$ is (Dirac-mode number)^2. On the L^4 lattice, the Dirac-mode number is $L^4\times N_c\times$ 4, which can be reduced to be $L^4\times N_c$, using the KS technique [3, 5], as mentioned in Sec.II-C. The actual number of the independent Dirac eigenvalue λ_n is about $L^4\times N_c/2$, due to the chiral property of $\not{D}$, i.e., pairwise appearance of $\pm\lambda_n$. Even for the projected operators, where the Dirac-mode space is restricted, the matrix is generally still huge. In addition, we have to deal with the product of the huge matrices $\langle m|\hat{U}_\mu|n\rangle$ in calculating the Wilson loop. Then, at present, we use a small-size lattice in the numerical calculation.

In this paper, we use SU(3) lattice QCD with the standard plaquette action at $\beta=5.6$ on 6^4 at the quenched level. The lattice QCD Monte Carlo calculation is done using the pseudo-heat-bath algorithm. The lattice spacing a is estimated as $a \simeq 0.25$fm, i.e., $a^{-1} \simeq 0.8$GeV, which leads to the string tension $\sigma \simeq 0.89$GeV/fm in the inter-quark potential. Then, the total volume is $V = (6a)^4 \simeq (1.5\text{fm})^4$, and the momentum cutoff is $\pi/a \simeq 2.5$GeV.

On the 6^4 lattice, the Dirac-mode number is $6^4 \times 3 \times 4 = 15,552$, which is reduced to be $6^4 \times 3 = 3,888$ using the KS technique. In fact, the KS Dirac operator $\eta_\mu D_\mu$ and the KS-reduced matrix element $\langle m|\hat{U}_\mu|n\rangle$ are expressed by $3,888 \times 3,888$ matrix. Considering the pairwise appearance of λ_n and $-\lambda_n$ apart from chiral zero modes, the actual number of the independent Dirac eigenvalue λ_n is reduce to be around $6^4 \times 3/2 = 1,944$.

To diagonalize the KS Dirac operator $\eta_\mu D_\mu$, we use LAPACK [30]. For the statistical error on the lattice data, we adopt the jackknife error estimate [5].

We show in Fig.1(a) the spectral density $\rho(\lambda)$ of the QCD Dirac operator $\not{D}$. The chiral property of $\not{D}$ leads to $\rho(-\lambda) = \rho(\lambda)$. Figure 1(b) is the IR-cut Dirac spectral density

$$\rho_{\rm IR}(\lambda) \equiv \rho(\lambda)\theta(|\lambda| - \Lambda_{\rm IR}) \tag{33}$$

with the IR-cutoff $\Lambda_{\rm IR} = 0.5a^{-1} \simeq 0.4$GeV.



FIG. 1: (a) The spectral density $\rho(\lambda)$ of the Dirac operator in lattice QCD at β=5.6 and 6^4. The negative-λ region is omitted, because of $\rho(-\lambda) = \rho(\lambda)$. The volume V is multiplied. (b) The IR-cut Dirac spectral density $\rho_{\rm IR}(\lambda) \equiv \rho(\lambda)\theta(|\lambda| - \Lambda_{\rm IR})$ with the IR-cutoff $\Lambda_{\rm IR} = 0.5a^{-1} \simeq 0.4$GeV.

Note that, using the eigenvalue λ_n, the quark condensate $\langle \bar{q}q \rangle$ is obtained as

$$\begin{aligned} \langle \bar{q}q \rangle &= {\rm Tr}\frac{1}{\not{D} + m} = \sum_n \frac{1}{i\lambda_n + m} \\ &= \sum_{\lambda_n > 0} \frac{2m}{\lambda_n^2 + m^2} + \frac{\nu}{m^2}, \end{aligned} \tag{34}$$

where ν is the total number of the zero mode of $\not{D}$. Here, apart from the zero mode, the eigenvalues appear as pairwise, which makes $\langle \bar{q}q \rangle$ real. (In lattice QCD, one has to take account of the doubler contribution, which can be regarded as flavor at the quenched level.) Then, in the presence of the IR cut $\Lambda_{\rm IR}$ for the Dirac eigen-mode, the quark condensate is obtained as

$$\langle \bar{q}q \rangle_{\Lambda_{\rm IR}} = \sum_{\lambda_n \ge \Lambda_{\rm IR}} \frac{2m}{\lambda_n^2 + m^2}. \tag{35}$$

We show in Fig.2 the lattice QCD result of the quark condensate $\langle \bar{q}q \rangle_{\Lambda_{\rm IR}}$ as the function of the current quark mass m in the presence of IR cut $\Lambda_{\rm IR}$.



FIG. 2: The lattice QCD result of the quark condensate $\langle \bar{q}q \rangle_{\Lambda_{\rm IR}}$ as the function of the current quark mass m in the presence of IR cut $\Lambda_{\rm IR} = 0.5, 1.0, 1.5[a^{-1}]$. The vertical axis is normalized by the original value of $\langle \bar{q}q \rangle$ without cut. A large reduction is found as $\langle \bar{q}q \rangle_{\Lambda_{\rm IR}}/\langle \bar{q}q \rangle \simeq 0.02$ for $\Lambda_{\rm IR} = 0.5a^{-1} \simeq 0.4$GeV around the physical region of $m \simeq 0.006a^{-1} \simeq 5$MeV.

By removing the low-lying Dirac modes, the chiral condensate $\langle \bar{q}q \rangle$ is largely reduced, reflecting the Banks-Casher relation. Actually, directly from lattice QCD calculation, we find a large reduction of the chiral condensate in the presence of the IR cut $\Lambda_{\rm IR} = 0.5a^{-1} \simeq 0.4$GeV,

$$\frac{\langle \bar{q}q \rangle_{\Lambda_{\rm IR}}}{\langle \bar{q}q \rangle} \simeq 0.02, \tag{36}$$

around the physical region of $m \simeq 0.006a^{-1} \simeq 5$MeV [31], as shown Fig.2.

Figure 3 shows the Dirac-mode projected inter-quark potential $V^P(R)$ after removing low-lying Dirac modes, which is obtained in lattice QCD with the IR-cut of $\rho_{\rm IR}(\lambda) \equiv \rho(\lambda)\theta(|\lambda| - \Lambda_{\rm IR})$ with the IR-cutoff $\Lambda_{\rm IR} = 0.5a^{-1}$. Remarkably, no significant change is observed



FIG. 3: The lattice QCD result of the inter-quark potential $V^P(R)$ after removing low-lying Dirac modes. The circle symbol denotes the potential obtained with the IR-cut of $\rho_{\rm IR}(\lambda) \equiv \rho(\lambda)\theta(|\lambda| - \Lambda_{\rm IR})$ with the IR-cutoff $\Lambda_{\rm IR} = 0.5a^{-1}$. The square symbol denotes the original inter-quark potential. The potential is almost unchanged even after removing the low-lying Dirac modes, apart from an irrelevant constant.

on the inter-quark potential besides an irrelevant constant, that is, the confining force is almost unchanged, even after removing the low-lying Dirac modes, which are responsible to chiral symmetry breaking [25]. This result indicates that one-to-one correspondence does not hold for confinement and chiral symmetry breaking in QCD.

We also investigate the UV-cut of Dirac modes in lattice QCD, using the UV-cut Dirac spectral density $\rho_{\rm UV}(\lambda) \equiv \rho(\lambda)\theta(\Lambda_{\rm UV} - |\lambda|)$ with the UV-cutoff $\Lambda_{\rm UV} = 2a^{-1} \simeq 1.6$GeV. In this case, unlike the IR cut, the chiral condensate is almost unchanged, and chiral symmetry breaking is almost kept. We show in Fig. 4 the UV-cut inter-quark potential, after removing the UV Dirac modes. We find that the confining force is almost unchanged by the UV-cut. This result seems consistent with the pioneering lattice study of Synatschke-Wipf-Langfeld [16]: they found that the confinement potential is almost reproduced only with low-lying Dirac modes, using the spectral sum of the Polyakov loop [32].



FIG. 4: (a) The UV-cut Dirac spectral density $\rho_{\rm UV}(\lambda) \equiv \rho(\lambda)\theta(\Lambda_{\rm UV} - |\lambda|)$ with the UV-cutoff $\Lambda_{\rm UV} = 2a^{-1} \simeq 1.6$GeV. (b) The UV-cut inter-quark potential (circle) after removing the UV Dirac modes. The potential is almost unchanged from the original one (square), apart from an irrelevant constant.

Furthermore, we examine "intermediate(IM)-cut", where a certain part of $\Lambda_1 < |\lambda_n| < \Lambda_2$ of Dirac modes is removed. Unfortunately, when the wide region of Dirac modes is removed, the statistical error becomes quite large for the Dirac-mode projected inter-quark potential. Here, we remove the IM Dirac modes of $0.5 - 0.8[a^{-1}]$, $0.8 - 1.0[a^{-1}]$, and $1.0 - 1.2[a^{-1}]$, respectively, and investigate the corresponding IM-cut inter-quark potential in each case, as shown in Fig.5. For each case, the confining force is found to be almost unchanged by the IM-cut.



FIG. 5: The left figures show the intermediate(IM)-cut Dirac spectral density $\rho_{\rm IM}(\lambda)$: the IM Dirac modes of $0.5-0.8[a^{-1}]$ (top), $0.8-1.0[a^{-1}]$ (middle), and $1.0-1.2[a^{-1}]$ (bottom) are cut. The right figures show the corresponding IM-cut inter-quark potential (circle) after removing the IM Dirac modes. For each case, the potential is almost unchanged from the original one (square), apart from an irrelevant constant.

Thus, from the above lattice QCD results, we conclude that there is no specific region of the Dirac modes responsible to the confinement in QCD, unlike the chiral symmetry breaking. Instead, we conjecture that the "seed" of confinement is distributed not only in low-lying Dirac modes but also in a wider region of the Dirac-mode space.

VI. SUMMARY AND DISCUSSIONS

We have developed a manifestly gauge-covariant expansion using the eigen-mode of the QCD Dirac operator $\not{D} = \gamma^\mu D^\mu$. With this method, we have performed a direct investigation of correspondence between confinement and chiral symmetry breaking in lattice QCD. As a remarkable fact, the confinement force is almost unchanged even after removing the low-lying Dirac modes, which are responsible to chiral symmetry breaking. This indicates that one-to-one correspondence does not hold for between confinement and chiral symmetry breaking in QCD.

Our strategy is to investigate the relation between the nonperturbative properties of QCD, by extracting or removing the essence of chiral symmetry breaking. This is similar to the demonstration of Abelian/monopole dominance [9, 12–14, 20–24, 34] or center/vortex dominance [15, 35, 36] for nonperturbative properties. However, while the previous scenario has been done in a specific gauge, our new method is manifestly gauge-invariant. In this analysis, we have carefully amputated only the "essence of chiral symmetry breaking" by cutting off the low-lying Dirac modes. Then, we have artificially realized the "confined but chiral restored situation" in QCD.

Recently, Lang and Schrock studied the hadron spectra after the cut of the low-lying Dirac modes [33]. Since the quark propagator is directly expressed with the Dirac operator $\not{D}$, the Dirac-mode projection is straightforward, and complicated projection procedure is not necessary in such studies. In their study, although the confinement was not checked, the appearance of hadronic spectra seems to suggest the existence of the confinement force, even after cutting the low-lying Dirac modes.

Next, we comment on the possible relation among confinement, chiral symmetry breaking, and monopoles in QCD. There is a close relation between confinement and chiral symmetry breaking through the monopoles in the MA gauge [12–14]. The monopole would be essential degrees of freedom for most nonperturbative QCD: confinement [21], chiral symmetry breaking [13, 14], and instantons [22]. In fact, removing the monopole would be "too fatal" for the nonperturbative properties, so that nonperturbative QCD phenomena are simultaneously lost by their cut. On the approximate coincidence of the critical temperatures of deconfinement and chiral restoration, a large change of monopoles may lead to both phase transitions [13], since the global connection of the monopole current seems to be largely changed around the QCD phase transition [23].

As for the recent finite-temperature QCD analysis, a lattice QCD group has reported a certain difference between the "critical temperatures" of deconfinement and chiral restoration, which are determined by the susceptibility peak of the Polyakov loop and chiral condensate, respectively [37]. This may be also an indirect evidence of "confinement $\neq$ chiral symmetry breaking" in QCD.

Next, we briefly discuss the role of low-lying Dirac modes in the viewpoint of instantons in QCD. The Dirac zero-mode associated with an instanton is localized around it [11]. However, the localized objects are hard to contribute to the large-distance phenomenon of confinement, although such low-lying Dirac modes contribute to chiral symmetry breaking. Recall that instantons contribute to chiral symmetry breaking, but do not directly lead to confinement [11]. Then, as a thought experiment, if only instantons can be carefully removed from the QCD vacuum, confinement properties would be almost unchanged, but the chiral condensate is largely reduced, and accordingly some low-lying Dirac modes disappear. Thus, in this case, confinement is almost unchanged, in spite of the large reduction of low-lying Dirac modes.

If the relation between confinement and chiral symmetry breaking is not one-to-one in QCD, richer phase structure is expected in QCD. For example, the phase transition point can be different between deconfinement and chiral restoration in the presence of strong electromagnetic fields, because of their nontrivial effect on chiral symmetry [38].

To obtain more definite conclusion, it is desired to perform larger-volume lattice QCD calculations and to cut various wider region of the Dirac modes, although it is quite difficult. It is also interesting to investigate the similar analysis at finite temperatures in lattice QCD. The full QCD calculation in this direction is also an interesting subject.

Acknowledgements

H.S. is supported in part by the Grant for Scientific Research [(C) No. 23540306, Priority Arias "New Hadrons" (E01:21105006)] and T.I is supported by Grant-in-Aid for JSPS Fellows (No.23-752). from the Ministry of Education, Culture, Science and Technology (MEXT) of Japan. This work is supported by the Global COE Program, "The Next Generation of Physics, Spun from Universality and Emergence". The lattice QCD calculations are done on NEC SX-8R at Osaka University.

---

[1] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122**, 345 (1961); ibid. **124** 246, (1961) .
[2] K.G. Wilson, Phys. Rev. D**10**, 2445 (1974).
[3] J.B. Kogut and L. Susskind, Phys. Rev. D**11**, 395 (1975).
[4] M. Creutz, Phys. Rev. Lett. **43**, 553 (1979); Phys. Rev. D**21**, 2308 (1980).
[5] H.J. Rothe, Lattice Gauge Theories, (World Scientific, 2005), and its references.
[6] M. Creutz, "Quarks, Gluons and Lattices" (Cambridge University Press, 1985); Acta Phys. Slov. **61**, 1 (2011), and its references.
[7] Y. Nambu, Phys. Rev. D**10**, 4262 (1974); G. 't Hooft, in "High Energy Physics" (1975); S. Mandelstam, Phys. Rept. **23**, 245 (1976).
[8] T. Banks and A. Casher, Nucl. Phys. B**169**, 103 (1980).
[9] G. 't Hooft, Nucl. Phys. B**190**, 455 (1981).
[10] J.M. Cornwall, Phys. Rev. D**26**, 1453 (1982).
[11] M. Shifman, Instantons in Gauge Theories, (World Sci-

entific, 1994), and its references.

[12] H. Suganuma, S. Sasaki, and H. Toki, Nucl. Phys. B**435**, 207 (1995); S. Sasaki, H. Suganuma, and H. Toki, Prog. Theor. Phys. **94**, 373 (1995); S. Umisedo, H. Suganuma, and H. Toki, Phys. Rev. D**57**, 1605 (1998).

[13] O. Miyamura, Phys. Lett. B**353**, 91 (1995).

[14] R.M. Woloshyn, Phys. Rev. D**51**, 6411 (1995).

[15] R. Hollwieser, M. Faber, J. Greensite, U.M. Heller, and S. Olejnik, Phys. Rev. D**78**, 054508 (2008).

[16] F. Synatschke, A. Wipf, and K. Langfeld, Phys. Rev. D**77**, 114018 (2008).

[17] J.M. Cornwall, Phys. Rev. D**83**, 076001 (2011).

[18] J.M. Cornwall, J. Papavassiliou, and D. Binosi, "The Pinch Technique and its Applications to Non-Abelian Gauge Theories", (Cambridge University Press, 2011) and its references.

[19] F. Karsch, Lect. Notes Phys. **583**, 209 (2002), and its references.

[20] A.S. Kronfeld, G. Schierholz, and U.-J. Wiese, Nucl. Phys. B**293**, 461 (1987); A.S. Kronfeld, M.L. Laursen, G. Schierholz, and U.-J. Wiese, Phys. Lett. B**198**, 516 (1987).

[21] J.D. Stack, S.D. Neiman, and R.J. Wensley, Phys. Rev. D**50**, 3399 (1994).

[22] H. Suganuma, A. Tanaka, S. Sasaki, and O. Miyamura, Nucl. Phys. **B** (Proc. Suppl.) **47**, 302 (1996).

[23] S.-I. Kitahara, Y. Matsubara, and T. Suzuki, Prog. Theor. Phys. **93**, 1 (1995).

[24] K. Amemiya and H. Suganuma, Phys. Rev. D**60**, 114509 (1999); H. Suganuma, K. Amemiya, H. Ichie, and A. Tanaka, Nucl. Phys. A**670**, 40 (2000).

[25] H. Suganuma, S. Gongyo, T. Iritani, and A. Yamamoto, PoS (**QCD-TNT II**), 044 (2011).

[26] A. Yamamoto and H. Suganuma, Phys. Rev. Lett. **101**, 241601 (2008) ; Phys. Rev. D**79**, 054504 (2009).

[27] A. Yamamoto and H. Suganuma, Phys. Rev. D**81**, 014506 (2010); PoS (**LAT2010**), 294 (2010); A. Yamamoto, Nucl. Phys. B**844**, 115 (2011).

[28] T. Iritani, H. Suganuma, and H. Iida, Phys. Rev. D**80**, 114505 (2009); T. Iritani and H. Suganuma, Phys. Rev. D**83**, 054502 (2011).

[29] M.F. Atiyah and I.M. Singer, Annals Math. **87**, 484 (1968).

[30] E. Anderson *et al.*, LAPACK Users' Guide, Third, (Society for Industrial and Applied Mathematics, 1999).

[31] K. Nakamura *et al.* (Particle Data Group), J. Phys. G**37**, 075021 (2010).

[32] C. Gattringer, Phys. Rev. Lett. **97** (2006) 032003; F. Bruckmann, C. Gattringer, and C. Hagen, Phys. Lett. B**647**, 56 (2007).

[33] C.B. Lang, M. Schrock, Phys. Rev. D**84**, 087704 (2011); PoS (**LAT2011**), 111 (2011).

[34] Z.F. Ezawa and A. Iwazaki, Phys. Rev. D**25**, 2681 (1982); Phys. Rev. D**26**, 631 (1982).

[35] L. Del Debbio, M. Faber, J. Greensite, and S. Olejnik, Phys. Rev. D**55** (1997) 2298;

[36] T.G. Kovacs and E.T. Tomboulis, Phys. Rev. D**57** (1998) 4054.

[37] Y. Aoki, Z. Fodor, S.D. Katz, and K.K. Szabo, Phys. Lett. B**643**, 46 (2006).

[38] H. Suganuma and T. Tatsumi, Ann. Phys. **208**, 470 (1991); Prog. Theor. Phys. **90**, 379 (1993).